

Paid Advertisement

SR/CL: May Day, 2009

Leitmotiv: "Under a spell so the wrong ones can't find it."

--R. Frost

שלאנקע שיפן

שלאנקע שיפן דרימלען אויפן געשוואלן גרינעם וואסער, שווארצע שאטנס שלאפן אויפן קאלטן הארץ פון וואסער. אלע ווינטן זיינען שטיל. כמארעס רוקן זיך געשפענסטיק אין דער נאכט דער שטומער. בלייך און רויק ווארט די ערד אויף בליץ און דונער. איך וועל זיין שטיל. --A. Margolin (Rosa L.)

The Loyal Order of Un-reconstructed Freaks in fraternal unity with The Fuzzy Furry Freak Brothers continue to upgrade Project Paperclip through the lens, *caeteris paribus*, of Los Alamos T.N.W.P. *vis-à-vis* ADW 5/1/2001, 3<sup>rd</sup>. Full report at the 479<sup>th</sup> Anniversary Celebration of the Confessio Augustana, Hotel California-Salem Hall, 1200 Hours. The Little Orphan Children have posted the Advance Proceedings steganographically (2<sup>nd</sup> Pass as per 5/1/1990, 1<sup>st</sup> L.) amongst the locallyokelpig-pondscum's *Senfgrube* of pictorial files. (Aesthetically amusing how they delight in showing so much fallow, sagging mortality—all of it soon to be smoke up the stack. Could they read, this open transmission would be securely published—or floated—in the cloacal *ambiance* of the aforesaid Treponema pallidum. The Hirelings, however, have given THE ORPHANAGE considerable pause with their latest gambit; transmit all future responses *via* the Sobol Sequence (*und nicht* Antonov/Saleev). Winthrop and Endecott have separately verified the Arbella's cybernetic manifest; Anne Bradstreet will take delivery *sub specie* 5/1/2007, 3<sup>rd</sup> L., on Trinity Sunday, 2009, Foster's Wharf—extrapolate hour and minute: 5/1/89, matrix [Romberg-Beleznay *cum* Thomas Bradwardine's (patented) *sub rosa* modification.]. Farragut has secured an additional vat of Full Doctrine to be received by Jenny Geddes at The Steelyard (*Einshof*) with The Revisionists seeing only the bustle of Cannon Street Station [EC4N6AP], even as history is made under their distracted gaze. The greatest *coup* since we went invisible—**LAUS DEO**!—has been against the Pelagian devils themselves—our only shape-shifting opponents: channel-rhodopsins mated with 4-ethylguaiacol. Since last year's Celebration of the Confessio, the first field tests of [non-flawed] *Rosumovi Univerzální Roboti* have demonstrated to (an admittedly preliminary, non-Bayesian)  $t=.02$ ,  $p<.05$ , that Pigs guzzling *vin rouge* of any grade, respond to the empirical exemplars. Their "will"—if they possessed such—is statistically immaterial when pitted against the repetitive imbibition coupled with THE ORPHANAGE's imaginal archetypes. Иван Петрович Павлов, you should be living at this hour. *"Et ad haec quis tam idoneus?"* **FRATRES, ORATE PRO NOBIS.**



Martín Lutero

IC XC  
NIKA


ology Porter-Silver color reaction or Nymberg method for  
nic steroids  
nal Information The Porter-Silver color reaction detects ster  
diacyclopropane group at carbon 17, including major glucocor  
etabolites. By preheating with a strong reducing agent, c  
ses the number of metabolites detected (ie, 17-ketogestrol  
s). Either 17-hydroxyprogesterone or 17-ketogenic steroid metabo  
can be used as an estimate of adrenal steroid production, g  
ealve values or in stimulation or suppression tests.  
xortosterone (DOC), corticosterone (compound B) and c  
lack a 17-hydroxyl group and are not detected in 17-KG and  
procedures.  
he  
per MA, Nieren U, Culler GB, et al, "Primary Steroid Adrenalocortical C  
high's Syndrome," Surgery, 1981, 111(6):1105-15.  
ck MP, Olfendick GA, Culler GB, et al, "Urea Free Cortisol in the High-C  
urethane Suppression Test for the Differential Diagnosis of the C  
drome," Ann Intern Med, 1982, 119(3):217.  
ces  
IL, Glass RH, and Kane HS, Clinical Gynecologic Endocrinology and And  
rology, Baltimore, MD: Williams & Wilkins, 1989.  
xyestroladiol see Estradiol, Unconjugated, Pregnancy, Blo  
s on page 126  
xyeticholanolone see 17-Ketosteroids Fractionation,  
n page 152  
xyindoleacetic Acid, Quantitative, Urine  
197  
Information  
rin on page 136  
Collection, 24-hour on page 30  
ms 5-HIAA, Quantitative, Urine  
to Serotonin, Metabolite  
it A serotonin metabolite, urinary excretion of 5-hydrox  
cetic acid (5-HIAA) occurs in the presence of carinoid line  
id tumors are enterochromaffin neoplasms containing an  
ar uptake and decarboxylation cells (APUDomas). They occur  
ile endocrine neoplasia syndrome (MEN) I or II but are often  
of those entities.  
Preparation Avoid bananas, avocados, chocolate, p  
nt, tomatoes, plantain, pineapples, melons, avocados, and  
ase, pheasant, tough egnip containing glyceryl guaiacate  
n (Baqasyn<sup>®</sup>, Anuprot<sup>®</sup>, neptegren, metocarbamol, m  
sonant, MAO inhibitors, mefenamic, methyllopho, pheno  
grose carcinoid tumors and syndromes, values >25 mg/24 ho  
n. Patients with renal disease may have falsely low 5-H  
n the urine. 5-HIAA is increased in untreated patients with m  
n, who have increased urinary tryptophan metabolites. Su  
a include those with colic disease, tropical sprue, Whipple  
a, stasis syndrome, and cystic fibrosis. It is increased in those  
intestinal obstruction and with some noncarcinoid (ie, c  
Poor correlation exists between 5-HIAA level and the c  
y of the carcinoid syndrome. It is used as a prognostic fac  
eas.  
ndications False elevated results if foods containing sero  
a producing metabolites that react with nitroacetylphor, a re  
this determination, are ingested within 2 days prior to collect  
ology Colonytely after extraction, spectrophotometric, gas chromat  
nity (GC), high performance liquid chromatography (HPLC), or page 147  
canza plasmation immunosay (PPAI) A carbon fiber elect  
as allows in vivo monitoring of 5-HIAA from small brain nucle  
nal Information 5-HIAA is the major urinary metabolite of seru  
a ubiquitous bioactive amine. Serotonin, and consequently, 5-H  
are produced in excess by most carcinoid tumors, especim  
um-Osmolality see Osmolality, Serum on page 172  
used

IG MONITORING/TOXICOLOGY/DRUGS OF ABU  
cretin Test see Gastrin, Serum on page 135  
cretin Test see Gastrin, Serum on page 135  
etabolites. By preheating with a strong reducing agent, c  
ses the number of metabolites detected (ie, 17-ketogestrol  
s). Either 17-hydroxyprogesterone or 17-ketogenic steroid metabo  
can be used as an estimate of adrenal steroid production, g  
ealve values or in stimulation or suppression tests.  
xortosterone (DOC), corticosterone (compound B) and c  
lack a 17-hydroxyl group and are not detected in 17-KG and  
procedures.  
he  
per MA, Nieren U, Culler GB, et al, "Primary Steroid Adrenalocortical C  
high's Syndrome," Surgery, 1981, 111(6):1105-15.  
ck MP, Olfendick GA, Culler GB, et al, "Urea Free Cortisol in the High-C  
urethane Suppression Test for the Differential Diagnosis of the C  
drome," Ann Intern Med, 1982, 119(3):217.  
ces  
IL, Glass RH, and Kane HS, Clinical Gynecologic Endocrinology and And  
rology, Baltimore, MD: Williams & Wilkins, 1989.  
xyestroladiol see Estradiol, Unconjugated, Pregnancy, Blo  
s on page 126  
xyeticholanolone see 17-Ketosteroids Fractionation,  
n page 152  
xyindoleacetic Acid, Quantitative, Urine  
197  
Information  
rin on page 136  
Collection, 24-hour on page 30  
ms 5-HIAA, Quantitative, Urine  
to Serotonin, Metabolite  
it A serotonin metabolite, urinary excretion of 5-hydrox  
cetic acid (5-HIAA) occurs in the presence of carinoid line  
id tumors are enterochromaffin neoplasms containing an  
ar uptake and decarboxylation cells (APUDomas). They occur  
ile endocrine neoplasia syndrome (MEN) I or II but are often  
of those entities.  
Preparation Avoid bananas, avocados, chocolate, p  
nt, tomatoes, plantain, pineapples, melons, avocados, and  
ase, pheasant, tough egnip containing glyceryl guaiacate  
n (Baqasyn<sup>®</sup>, Anuprot<sup>®</sup>, neptegren, metocarbamol, m  
sonant, MAO inhibitors, mefenamic, methyllopho, pheno  
grose carcinoid tumors and syndromes, values >25 mg/24 ho  
n. Patients with renal disease may have falsely low 5-H  
n the urine. 5-HIAA is increased in untreated patients with m  
n, who have increased urinary tryptophan metabolites. Su  
a include those with colic disease, tropical sprue, Whipple  
a, stasis syndrome, and cystic fibrosis. It is increased in those  
intestinal obstruction and with some noncarcinoid (ie, c  
Poor correlation exists between 5-HIAA level and the c  
y of the carcinoid syndrome. It is used as a prognostic fac  
eas.  
ndications False elevated results if foods containing sero  
a producing metabolites that react with nitroacetylphor, a re  
this determination, are ingested within 2 days prior to collect  
ology Colonytely after extraction, spectrophotometric, gas chromat  
nity (GC), high performance liquid chromatography (HPLC), or page 147  
canza plasmation immunosay (PPAI) A carbon fiber elect  
as allows in vivo monitoring of 5-HIAA from small brain nucle  
nal Information 5-HIAA is the major urinary metabolite of seru  
a ubiquitous bioactive amine. Serotonin, and consequently, 5-H  
are produced in excess by most carcinoid tumors, especim  
um-Osmolality see Osmolality, Serum on page 172  
used

that, blood levels cannot be well predicted from doing sch  
Furthermore, this drug has a narrow therapeutic window and sig  
cidity at levels above that range.  
ions Results are method dependent - some measure m  
dies as well as parent drug. Single assays are not as inform  
eas over time.  
ology-Radiimmunoassay (RIA), high performance liquid  
nal Information  
-HIT: 5-Hydroxytryptamine, Blood,  
Applies to Serotonin, Cerebrospinal Fluid  
Abstract Sacrofen (5-Hydroxytryptamine [5-HT]) is synthe  
typtophan. Sacrofen's major metabolite, 5-HIAA, is meas  
Commonly than 5-HT (parent compound). This test is rarely upo  
to likely to be administered to a transplant recipient. Include an  
different Preparation Monoamine oxidase inhibitor drugs  
discontinued for at least 1 week prior to sampling, since they  
increase the level of serotonin. Avoid application of radiod  
scars) before collection of specimen if RIA is used for as  
methods require a low iodine diet for several days. Avoid toxic  
sacacchi, bananas, kiwifruit, pineapple, walnuts, and nut  
Specimen Whole blood, cerebrospinal fluid  
Container Tube with EDTA, sometimes with ascorbic acid,  
laboratory if the assay is available at all.  
Collection Draw in chilled tubes. Keep on ice.  
Storage Instructions Place whole blood in plastic bottle co  
mg EDTA and 75 mg ascorbic acid. Freeze within 4 hours of  
Stable 7 days at -20°C.  
Reference Range 10-30 µg/dL (SI: 570-1700 nmol/L).  
among laboratories and are method dependent. In serum, 5-HIAA  
hydroxytryptamine levels in females were about 1.3-fold that increased concentrations of cyclosporine include telomeres  
By RIA, a study provided ranges: male 7-12 µg/dL (SI: 380-600 nmol/L), female 0-16 µg/dL (SI: 520-300 nmol/L).  
which enhances the potential toxicity of cyclosporine, and i  
to likely to be administered to a transplant recipient. Include an  
side antibiotics, cephalosporins, trimethoprim-sulfam, aspirin  
dow, valproic acid, and fentanyl. Agents which raise cy  
Use levels by decreasing biotransformation include methylphen  
Methods of cyclosporine assay include immunoassay, HPLC, and  
Abstract This drug is a sedative-hypnotic of the benzodiaz  
with a rather wide therapeutic window.  
Specimen Serum  
Container Red top tube  
Reference Range Therapeutic: 0.4 ng/mL (SI: 0.9 nmol/L)  
N-desethylflurazepam: 20-110 ng/mL (SI: 43-240 nmol/L)  
N-desethylflurazepam: 20-110 ng/mL (SI: 43-240 nmol/L)  
Possible Panic Range Toxic: 200 ng/mL (SI: 500 nmol/L)  
Use Monitor therapeutic drug level (nearly), toxicity assessme  
Methodology Thin-layer chromatography (TLC), high perform  
chromatography (HPLC)  
Additional Information  
• Half-life: parent drug: 3-6 hours; metabolite (N-desethyl)  
50-100 hours  
• Volume of distribution: 15-30 L/kg  
• Protein binding: 98% to 99%  
Most common side effect is daytime drowsiness. The m  
metabolite is N-1-hydroxyethyl flurazepam.  
References  
Griesbach DJ, Hemsch JS, Engelhardt N, et al, Pharmacokinetic Di  
Dynamic Differences Among Three Benzodiazepine Hypnotics, Fluraz  
epam, and Temazepam," Arch Gen Psychiatry, 1989, 46(4):338-52.  
Riles A, "Benzodiazepine Hypnotics and Anesthetics," Hosp Pract (C  
25 Sept 9:7-12.  
lex<sup>®</sup> see Methicarbamol on page 560  
onlex<sup>®</sup> see Flurazepam on previous page.  
ortunam<sup>®</sup> see Haloperidol on next page.  
ree Base see Cocaine (Cocaine Metabolite), Qualitative,  
page 543<sup>®</sup>  
ree Phenylephrine see Phenylephrine, Free on page 570  
urazepam<sup>®</sup> see Amitriptyline B on page 535  
abapentin  
CPT 82239  
Abstract Gabapentin is a GABAergic anticonvulsant effect  
and secondarily generalized seizures. Its lack of interactions  
anticonvulsants makes it useful as add-on therapy in refract  
epilepsy.  
Specimen Plasma  
Container Green top (heparin) tube  
Sampling Time Peak: 1.5-3 hours after a 200 mg dose  
Storage Instructions Refrigerate  
Reference Range Peak plasma level: 2 µg/mL, 1.5-3 hours  
mg dose  
Additional Information  
• Half-life: 5-7 hours  
• Volume of distribution: 1.0 L/kg  
• Protein binding: <3%  
Therapeutic monitoring is not required with gabapentin. As  
gabapentin is dose related, with less absorption at doses  
resulting in less toxicity after large ingestions. Fully pers  
absorbed after an oral dose of gabapentin, with a peak conc  
hours after administration. A distinct advantage of gabapentin  
cleared today by the kidneys so it is not subject to and dose  
changes in hepatic enzyme activity. Gabapentin has not be  
to interact with any other antiepileptic. Dosage adjustme



"Las cartas de navegación ya no eran secretos bien guardados, hasta los malditos ingleses las poseían." --I. Allende, 10:0-06-116156X, pp. 58-59.



Correlate these MAC generators-PIGS.